Albania Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)





Albania has a population of 1.20 million women ages 15 years and older who are at risk of developing cervical cancer. Current estimates indicate that every year 133 women are diagnosed with cervical cancer and 74 die from the disease.

Cervical cancer ranks as the 5th most frequent cancer among women in Albania and the 2nd most frequent cancer among women between 15 and 44 years of age. Data is not yet available on the HPV burden in the general population of Albania. However, in Southern Europe, the region Albania belongs to, about 3.8% of women in the general population are estimated to harbour cervical HPV-16/18 infection at a given time, and 68.0% of invasive cervical cancers are attributed to HPVs 16 or 18.

Table 1. Crude incidence rates of HPV-related cancers

	Male	Female
Cervical cancer	-	9.41
Anal cancer	0.20	0.07
Vulva cancer	-	1.70
Vaginal cancer	-	0.57
Penile cancer	0.68	-
Oropharyngeal cancer	1.09	0.85
Oral cavity cancer	3.96	3.40
Laryngeal cancer	8.53	1.56

Table 2. Burden of cervical cancer

	Incidence	Mortality
Annual number of new cases/deaths	133	74
Crude rate	9.41	5.24
Age-standarized rate	6.64	3.25
Cumulative risk 0-74 years (%)	0.71	0.38
Ranking of cervical cancer (all years)	5th	6th
Ranking of cervical cancer (15-44 years)	2nd	6th

Table 3. Burden of cervical HPV infection Albania

	No. Tested	% (95% CI)
HPV 16/18 prevalence:		
Normal cytology	-	-
Low-grade cervical lesions	-	-
High-grade cervical lesions	-	-
Cervical cancer	-	-

Figure 1. Comparison of the ten most frequent HPV oncogenic types in Albania among women with and without cervical lesions

No data available

Albania Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)



II. Complementary data on cervical cancer prevention

 Table 4. Factors contributing to cervical cancer (co-factors)

Smoking prevalence (%) [95% UI], women	7.5 [4.4-10.8]
Total fertility rate (live births per women)	1.7
Hormonal contraception use (%)	0.60
HIV prevalence (%) [95% UI] (15-49 years)	- [—]

Table 5. Sexual behaviour

MEN	
Percentage of 15-year-old who have had sexual intercourse	39.0
Range of median age at first sexual intercourse	17.0 - 23.4
WOMEN	
Percentage of 15-year-old who have had sexual intercourse	2.0
Range of median age at first sexual intercourse	20.0 - 21.3


Table 6. HPV vaccine introduction

Females	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-
Males	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-

Table 7. Cervical screening practices and recommendations

Incidations	
Existence of official national rec-	Yes
ommendations	
Starting year of recommendations	2019
Active invitation to screening	Yes
Screening ages (years), pri- mary screening test used, and screening interval or frequency of	40-50 (HPV test, 5 years)
screenings	

Figure 2. Estimated coverage of cervical cancer screening in Albania*



* Estimated coverage and 95% confidence interval in 2019

Contact information:

ICO/IARC HPV Information Centre Institut Català d'Oncologia Avda. Gran Via de l'Hospitalet, 199-203 08908 L'Hospitalet de Llobregat (Barcelona, Spain) info@hpvcentre.net

www.hpvcentre.net