Bangladesh

HPV INFORMATION CENTRE

Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)

I. Key data on HPV and HPV-related cancers



Bangladesh has a population of 64.0 million women ages 15 years and older who are at risk of developing cervical cancer. Current estimates indicate that every year 8268 women are diagnosed with cervical cancer and 4971 die from the disease. Cervical cancer ranks as the

2nd most frequent cancer among women in Bangladesh and the 2nd most frequent cancer among women between 15 and 44 years of age. Data is not yet available on the HPV burden in the general population of Bangladesh. However, in Southern Asia, the region Bangladesh belongs to, about 4.4% of women in the general population are estimated to harbour cervical HPV-16/18 infection at a given time, and 80.3% of invasive cervical cancers are attributed to HPVs 16 or 18.

Table 1. Crude incidence rates of HPV-related cancers

	Male	Female
Cervical cancer	-	10.2
Anal cancer	0.33	0.13
Vulva cancer	-	0.32
Vaginal cancer	-	0.32
Penile cancer	0.77	-
Oropharyngeal cancer	3.92	0.72
Oral cavity cancer	11.2	5.68
Laryngeal cancer	5.49	0.86

Table 2. Burden of cervical cancer

	Incidence	Mortality
Annual number of new cases/deaths	8268	4971
Crude rate	10.2	6.10
Age-standarized rate	10.6	6.67
Cumulative risk 0-74 years (%)	1.16	0.76
Ranking of cervical cancer (all years)	2nd	3rd
Ranking of cervical cancer (15-44 years)	2nd	2nd

Table 3. Burden of cervical HPV infection Bangladesh

	No. Tested	% (95% CI)
HPV 16/18 prevalence:		
Normal cytology	-	-
Low-grade cervical lesions	13	23.1 (8.2-50.3)
High-grade cervical lesions	-	-
Cervical cancer	-	-

Figure 1. Comparison of the ten most frequent HPV oncogenic types in Bangladesh among women with and without cervical lesions

No data available

Bangladesh



Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)

II. Complementary data on cervical cancer prevention

Table 4. Factors contributing to cervical cancer (cofactors)

Smoking prevalence (%) [95% UI], women	0.9 [0.5-1.2]
Total fertility rate (live births per women)	2.1
Hormonal contraception use (%)	27
HIV prevalence (%) [95% UI] (15-49 years)<	<0.1 [<0.1 -<0.1]

Table 5. Sexual behaviour

TUDIO OI COMUNI DOMATICAL	
MEN	
Percentage of 15-year-old who have had sexual intercourse	6.9
Range of median age at first sexual intercourse	
WOMEN	
Percentage of 15-year-old who have had sexual intercourse	15.0
Range of median age at first sexual intercourse	15.8 - 16.2


Table 6. HPV vaccine introduction

Females	_
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-
Males	_
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-

Table 7. Cervical screening practices and recommendations

Existence of official national recommendations	Yes
Starting year of recommendations	2018
Active invitation to screening	No
Screening ages (years), primary screening test used, and screening interval or frequency of screenings	30-60 (VIA, 5 years)

Figure 2. Estimated coverage of cervical cancer screening in Bangladesh*



^{*} Estimated coverage and 95% confidence interval in 2019

Contact information:

ICO/IARC HPV Information Centre

Institut Català d'Oncologia

Avda. Gran Via de l'Hospitalet, 199-203

08908 L'Hospitalet de Llobregat (Barcelona, Spain)

info@hpvcentre.net

www.hpvcentre.net