

Bahrain

Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)



I. Key data on HPV and HPV-related cancers



Bahrain has a population of 410,669 women ages 15 years and older who are at risk of developing cervical cancer. Current estimates indicate that every year 21 women are diagnosed with cervical cancer and 12 die from the disease. Cervical cancer ranks as the 8th

most frequent cancer among women in Bahrain and the 8th most frequent cancer among women between 15 and 44 years of age. Data is not yet available on the HPV burden in the general population of Bahrain. However, in Western Asia, the region Bahrain belongs to, about 2.5% of women in the general population are estimated to harbour cervical HPV-16/18 infection at a given time, and 72.4% of invasive cervical cancers are attributed to HPVs 16 or 18.

Table 1. Crude incidence rates of HPV-related cancers

	Male	Female
Cervical cancer	-	3.49
Anal cancer	0.27	0
Vulva cancer	-	0.17
Vaginal cancer	-	0
Penile cancer	0	-
Oropharyngeal cancer	0.09	0
Oral cavity cancer	1.27	0.67
Laryngeal cancer	1.09	0


Table 2. Burden of cervical cancer

	Incidence	Mortality
Annual number of new cases/deaths	21	12
Crude rate	3.49	2.00
Age-standardized rate	3.92	2.40
Cumulative risk 0-74 years (%)	0.48	0.31
Ranking of cervical cancer (all years)	8th	10th
Ranking of cervical cancer (15-44 years)	8th	7th

Table 3. Burden of cervical HPV infection Bahrain

	No. Tested	% (95% CI)
HPV 16/18 prevalence:		
Normal cytology	91	2.2 (0.6-7.7)
Low-grade cervical lesions	-	-
High-grade cervical lesions	-	-
Cervical cancer	-	-

Figure 1. Comparison of the ten most frequent HPV oncogenic types in Bahrain among women with and without cervical lesions



For data sources and further data on HPV infection in cancer sites other than the cervix, please refer to the specific country full report.

Bahrain

Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)

II. Complementary data on cervical cancer prevention

Table 4. Factors contributing to cervical cancer (co-factors)

Smoking prevalence (%) [95% UI], women	5.4 [2.7-8.4]
Total fertility rate (live births per women)	2.0
Hormonal contraception use (%)	10.9
HIV prevalence (%) [95% UI] (15-49 years)	- [—]

Table 5. Sexual behaviour

MEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	-
WOMEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	-


Table 6. HPV vaccine introduction

Females	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-
Males	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-

Table 7. Cervical screening practices and recommendations

Existence of official national recommendations	Yes
Starting year of recommendations	Unk
Active invitation to screening	No
Screening ages (years), primary screening test used, and screening interval or frequency of screenings	30-65 (cytology, 3 years)

Figure 2. Estimated coverage of cervical cancer screening in Bahrain*



* Estimated coverage and 95% confidence interval in 2019

Contact information:

ICO/IARC HPV Information Centre
 Institut Català d'Oncologia
 Avda. Gran Via de l'Hospitalet, 199-203
 08908 L'Hospitalet de Llobregat (Barcelona, Spain)
 info@hpvcentre.net
www.hpvcentre.net