


Belarus

Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)



I. Key data on HPV and HPV-related cancers



Belarus has a population of 4.36 million women ages 15 years and older who are at risk of developing cervical cancer. Current estimates indicate that every year 835 women are diagnosed with cervical cancer and 358 die from the disease.

Cervical cancer ranks as the 7th most frequent cancer among women in Belarus and the 3rd most frequent cancer among women between 15 and 44 years of age. About 7.1% of women in the general population are estimated to harbour cervical HPV-16/18 infection at a given time, and 65.4% of invasive cervical cancers are attributed to HPVs 16 or 18.

Table 1. Crude incidence rates of HPV-related cancers

	Male	Female
Cervical cancer	-	16.5
Anal cancer	0.80	1.15
Vulva cancer	-	4.75
Vaginal cancer	-	0.55
Penile cancer	1.25	-
Oropharyngeal cancer	12.1	0.55
Oral cavity cancer	12.4	2.79
Laryngeal cancer	14.2	0.38


Table 2. Burden of cervical cancer

	Incidence	Mortality
Annual number of new cases/deaths	835	358
Crude rate	16.5	7.09
Age-standardized rate	11.4	4.20
Cumulative risk 0-74 years (%)	1.12	0.44
Ranking of cervical cancer (all years)	7th	8th
Ranking of cervical cancer (15-44 years)	3rd	3rd

Table 3. Burden of cervical HPV infection Belarus

	No. Tested	% (95% CI)
HPV 16/18 prevalence:		
Normal cytology	322	7.1 (4.8-10.5)
Low-grade cervical lesions	94	35.1 (26.2-45.2)
High-grade cervical lesions	91	56.0 (45.8-65.8)
Cervical cancer	26	65.4 (46.2-80.6)

Figure 1. Comparison of the ten most frequent HPV oncogenic types in Belarus among women with and without cervical lesions



Belarus

Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)

II. Complementary data on cervical cancer prevention

Table 4. Factors contributing to cervical cancer (co-factors)

Smoking prevalence (%) [95% UI], women	9 [6.8-11.4]
Total fertility rate (live births per women)	1.7
Hormonal contraception use (%)	10.3
HIV prevalence (%) [95% UI] (15-49 years)	0.4 [0.3-0.5]

Table 5. Sexual behaviour

MEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	-
WOMEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	18.4 - 19.6


Table 6. HPV vaccine introduction

Females	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-
Males	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-

Table 7. Cervical screening practices and recommendations

Existence of official national recommendations	Yes
Starting year of recommendations	2019
Active invitation to screening	No
Screening ages (years), primary screening test used, and screening interval or frequency of screenings	>=18 (cytology, 2 years)

Figure 2. Estimated coverage of cervical cancer screening in Belarus*



* Estimated coverage and 95% confidence interval in 2019

Contact information:

ICO/IARC HPV Information Centre
 Institut Català d'Oncologia
 Avda. Gran Via de l'Hospitalet, 199-203
 08908 L'Hospitalet de Llobregat (Barcelona, Spain)
info@hpvcentre.net
www.hpvcentre.net