


Indonesia

Human Papillomavirus and Related Cancers, Fact Sheet 2021 (2021-10-22)

I. Key data on HPV and HPV-related cancers



Indonesia has a population of 99.8 million women ages 15 years and older who are at risk of developing cervical cancer. Current estimates indicate that every year 36633 women are diagnosed with cervical cancer and 21003 die from the disease. Cervical cancer ranks as

the 2nd most frequent cancer among women in Indonesia and the 2nd most frequent cancer among women between 15 and 44 years of age. About 4.0% of women in the general population are estimated to harbour cervical HPV-16/18 infection at a given time, and 87.0% of invasive cervical cancers are attributed to HPVs 16 or 18.

Table 1. Crude incidence rates of HPV-related cancers

	Male	Female
Cervical cancer	-	27.0
Anal cancer	0.36	0.20
Vulva cancer	-	0.98
Vaginal cancer	-	0.35
Penile cancer	0.74	-
Oropharyngeal cancer	0.76	0.32
Oral cavity cancer	2.58	1.64
Laryngeal cancer	2.48	0.18


Table 2. Burden of cervical cancer

	Incidence	Mortality
Annual number of new cases/deaths	36633	21003
Crude rate	27.0	15.5
Age-standardized rate	24.4	14.4
Cumulative risk 0-74 years (%)	2.69	1.73
Ranking of cervical cancer (all years)	2nd	2nd
Ranking of cervical cancer (15-44 years)	2nd	2nd

Table 3. Burden of cervical HPV infection Indonesia

	No. Tested	% (95% CI)
HPV 16/18 prevalence:		
Normal cytology	200	4.0 (2.0-7.7)
Low-grade cervical lesions	-	-
High-grade cervical lesions	-	-
Cervical cancer	230	87.0 (82.0-90.7)

Figure 1. Comparison of the ten most frequent HPV oncogenic types in Indonesia among women with and without cervical lesions



Indonesia

Human Papillomavirus and Related Cancers, Fact Sheet 2021 (2021-10-22)



II. Complementary data on cervical cancer prevention

Table 4. Factors contributing to cervical cancer (co-factors)

Smoking prevalence (%) [95% UI], women	2.6 [1.7-3.4]
Total fertility rate (live births per women)	2.3
Hormonal contraception use (%)	12.3
HIV prevalence (%) [95% UI] (15-49 years)	0.3 [0.3-0.4]

Table 5. Sexual behaviour

MEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	23.1 - 24.6
WOMEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	19.0 - 21.8

Table 6. HPV vaccine introduction in females

HPV vaccination programme	Partial introduction
Year of introduction	2017
HPV coverage – first dose (%) (estimation year 2019)	6
HPV coverage – last dose (%) (estimation year 2019)	6
HPV coverage – first dose (%) (estimation year 2020)	8
HPV coverage – last dose (%) (estimation year 2020)	8

Table 7. Cervical screening practices and recommendations

Cervical cancer screening coverage, % (age and screening interval, reference)	24.4% (All women aged 30-50 screened every 5y, Kim 2013)
Screening ages (years)	30-50
Screening interval (years) or frequency of screens	5 years

Figure 2. Estimated coverage of cervical cancer screening in Indonesia, by age and study

No data available

^a Proportion of women in the total sample of the mentioned age range in the country or region that reported having a Pap smear during a given time period (e.g., last year, last 2, 3, 5 years or ever).

Contact information:

ICO/IARC HPV Information Centre
 Institut Català d'Oncologia
 Avda. Gran Via de l'Hospitalet, 199-203
 08908 L'Hospitalet de Llobregat (Barcelona, Spain)
 info@hpvcentre.net
www.hpvcentre.net