


Kenya

Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)



I. Key data on HPV and HPV-related cancers



Kenya has a population of 16.8 million women ages 15 years and older who are at risk of developing cervical cancer. Current estimates indicate that every year 5236 women are diagnosed with cervical cancer and 3211 die from the disease. Cervical cancer ranks as the 2nd most frequent cancer among women in Kenya and the 2nd most frequent cancer among women between 15 and

44 years of age. About 9.1% of women in the general population are estimated to harbour cervical HPV-16/18 infection at a given time, and 63.1% of invasive cervical cancers are attributed to HPVs 16 or 18.

Table 1. Crude incidence rates of HPV-related cancers

	Male	Female
Cervical cancer	-	19.4
Anal cancer	0.22	0.54
Vulva cancer	-	0.58
Vaginal cancer	-	0.24
Penile cancer	0.18	-
Oropharyngeal cancer	0.18	0.07
Oral cavity cancer	1.46	1.36
Laryngeal cancer	1.26	0.27


Table 2. Burden of cervical cancer

	Incidence	Mortality
Annual number of new cases/deaths	5236	3211
Crude rate	19.4	11.9
Age-standardized rate	31.3	20.6
Cumulative risk 0-74 years (%)	3.60	2.50
Ranking of cervical cancer (all years)	2nd	1st
Ranking of cervical cancer (15-44 years)	2nd	2nd

Table 3. Burden of cervical HPV infection Kenya

	No. Tested	% (95% CI)
HPV 16/18 prevalence:		
Normal cytology	823	9.1 (7.3-11.3)
Low-grade cervical lesions	42	21.4 (11.7-35.9)
High-grade cervical lesions	20	45.0 (25.8-65.8)
Cervical cancer	233	63.1 (56.7-69.0)

Figure 1. Comparison of the ten most frequent HPV oncogenic types in Kenya among women with and without cervical lesions



Kenya

Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)



II. Complementary data on cervical cancer prevention

Table 4. Factors contributing to cervical cancer (co-factors)

Smoking prevalence (%) [95% UI], women	1 [0.7-1.4]
Total fertility rate (live births per women)	3.8
Hormonal contraception use (%)	4.92
HIV prevalence (%) [95% UI] (15-49 years)	6.1 [5-7.2]

Table 5. Sexual behaviour

MEN	
Percentage of 15-year-old who have had sexual intercourse	19.6
Range of median age at first sexual intercourse	16.7 - 17.6
WOMEN	
Percentage of 15-year-old who have had sexual intercourse	10.7
Range of median age at first sexual intercourse	16.7 - 18.2


Table 6. HPV vaccine introduction

Females	
HPV vaccination programme	Introduced
Year of introduction	2019
Year of estimation	2021
HPV coverage – first dose (%)	29
HPV coverage – last dose (%)	44
Males	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-

Table 7. Cervical screening practices and recommendations

Existence of official national recommendations	Yes
Starting year of recommendations	2018
Active invitation to screening	No
Screening ages (years), primary screening test used, and screening interval or frequency of years)	25-49 (VIA, 5 years); 25-30 (cytology, 5 years); 30-49 (HPV test, 5 years)

Figure 2. Estimated coverage of cervical cancer screening in Kenya*



* Estimated coverage and 95% confidence interval in 2019

Contact information:

ICO/IARC HPV Information Centre
 Institut Català d'Oncologia
 Avda. Gran Via de l'Hospitalet, 199-203
 08908 L'Hospitalet de Llobregat (Barcelona, Spain)
 info@hpvcentre.net
www.hpvcentre.net