


Libya

Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)



I. Key data on HPV and HPV-related cancers



Libya has a population of 2.40 million women ages 15 years and older who are at risk of developing cervical cancer. Current estimates indicate that every year 240 women are diagnosed with cervical cancer and 141 die from the disease. Cervical cancer ranks as the 3rd most frequent cancer among women in Libya and the 7th most frequent cancer among women between 15 and 44

years of age. Data is not yet available on the HPV burden in the general population of Libya. However, in Northern Africa, the region Libya belongs to, about 3.0% of women in the general population are estimated to harbour cervical HPV-16/18 infection at a given time, and 78.9% of invasive cervical cancers are attributed to HPVs 16 or 18.

Table 1. Crude incidence rates of HPV-related cancers

	Male	Female
Cervical cancer	-	7.05
Anal cancer	0.29	0.18
Vulva cancer	-	0.38
Vaginal cancer	-	0.24
Penile cancer	0	-
Oropharyngeal cancer	0.12	0.06
Oral cavity cancer	0.86	0.88
Laryngeal cancer	4.15	0.47

Table 2. Burden of cervical cancer

	Incidence	Mortality
Annual number of new cases/deaths	240	141
Crude rate	7.05	4.14
Age-standardized rate	7.70	4.88
Cumulative risk 0-74 years (%)	0.89	0.58
Ranking of cervical cancer (all years)	3rd	3rd
Ranking of cervical cancer (15-44 years)	7th	10th

Table 3. Burden of cervical HPV infection Libya

	No. Tested	% (95% CI)
HPV 16/18 prevalence:		
Normal cytology	-	-
Low-grade cervical lesions	-	-
High-grade cervical lesions	-	-
Cervical cancer	-	-

Figure 1. Comparison of the ten most frequent HPV oncogenic types in Libya among women with and without cervical lesions



Libya

Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)



II. Complementary data on cervical cancer prevention

Table 4. Factors contributing to cervical cancer (co-factors)

Smoking prevalence (%) [95% UI], women	- [—]
Total fertility rate (live births per women)	2.2
Hormonal contraception use (%)	8.20
HIV prevalence (%) [95% UI] (15-49 years)	0.1 [0.1-0.1]

Table 5. Sexual behaviour

MEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	-
WOMEN	
Percentage of 15-year-old who have had sexual intercourse	-
Range of median age at first sexual intercourse	-


Table 6. HPV vaccine introduction

Females	
HPV vaccination programme	Introduced
Year of introduction	2013
Year of estimation	2021
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-
Males	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-

Table 7. Cervical screening practices and recommendations

Existence of official national recommendations	No
Starting year of recommendations	-
Active invitation to screening	-
Screening ages (years), primary screening test used, and screening interval or frequency of screenings	-

Figure 2. Estimated coverage of cervical cancer screening in Libya*



* Estimated coverage and 95% confidence interval in 2019

Contact information:

ICO/IARC HPV Information Centre
 Institut Català d'Oncologia
 Avda. Gran Via de l'Hospitalet, 199-203
 08908 L'Hospitalet de Llobregat (Barcelona, Spain)
 info@hpvcentre.net
www.hpvcentre.net