Qatar



Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)

I. Key data on HPV and HPV-related cancers



Qatar has a population of 527,079 women ages 15 years and older who are at risk of developing cervical cancer. Current estimates indicate that every year 23 women are diagnosed with cervical cancer and 11 die from the disease. Cervical cancer ranks as the 5th

most frequent cancer among women in Qatar and the 3rd most frequent cancer among women between 15 and 44 years of age. Data is not yet available on the HPV burden in the general population of Qatar. However, in Western Asia, the region Qatar belongs to, about 2.5% of women in the general population are estimated to harbour cervical HPV-16/18 infection at a given time, and 72.4% of invasive cervical cancers are attributed to HPVs 16 or 18.

Table 1. Crude incidence rates of HPV-related cancers

	Male	Female
Cervical cancer	-	3.21
Anal cancer	0.18	0.84
Vulva cancer	-	0
Vaginal cancer	-	0
Penile cancer	0.05	-
Oropharyngeal cancer	0.05	0
Oral cavity cancer	1.15	0.70
Laryngeal cancer	0.83	0

Table 2. Burden of cervical cancer

	Incidence	Mortality
Annual number of new cases/deaths	23	11
Crude rate	3.21	1.54
Age-standarized rate	4.13	2.91
Cumulative risk 0-74 years (%)	0.53	0.36
Ranking of cervical cancer (all years)	5th	7th
Ranking of cervical cancer (15-44 years)	3rd	4th

Table 3. Burden of cervical HPV infection Qatar

	No. Tested	% (95% CI)
HPV 16/18 prevalence:		
Normal cytology	-	-
Low-grade cervical lesions	-	-
High-grade cervical lesions	-	-
Cervical cancer	-	-

Figure 1. Comparison of the ten most frequent HPV oncogenic types in Qatar among women with and without cervical lesions

No data available

Qatar



Human Papillomavirus and Related Cancers, Fact Sheet 2023 (2023-03-10)

II. Complementary data on cervical cancer prevention

Table 4. Factors contributing to cervical cancer (cofactors)

1401010)	
Smoking prevalence (%) [95% UI], women	0.9 [0.4-1.4]
Total fertility rate (live births per women)	1.9
Hormonal contraception use (%)	13.3
HIV prevalence (%) [95% UI] (15-49 years)	- [—]

Table 5 Sexual behaviour

lat	ole 5. Sexual behaviour	
MEI	N	
	Percentage of 15-year-old who have had sexual intercourse	-
	Range of median age at first sexual intercourse	-
WO	MEN	
	Percentage of 15-year-old who have had sexual intercourse	-
	Range of median age at first sexual intercourse	-


Table 6. HPV vaccine introduction

Females	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage – first dose (%)	-
HPV coverage – last dose (%)	-
Males	
HPV vaccination programme	Not Available/Not Introduced
Year of introduction	-
Year of estimation	-
HPV coverage - first dose (%)	-
HPV coverage – last dose (%)	-

Table 7. Cervical screening practices and recommendations

Existence of official national rec-	Yes
ommendations	
Starting year of recommendations	2012
Active invitation to screening	No
Screening ages (years), primary screening test used, and screening interval or frequency of screenings	21-49 (cytology, 3 years); 50-65 (cytology, 5 years)

Figure 2. Estimated coverage of cervical cancer screening in Qatar*



^{*} Estimated coverage and 95% confidence interval in 2019

Contact information:

ICO/IARC HPV Information Centre

Institut Català d'Oncologia

Avda. Gran Via de l'Hospitalet, 199-203

08908 L'Hospitalet de Llobregat (Barcelona, Spain)

info@hpvcentre.net

www.hpvcentre.net